

Gavin Newsom Mayor Communicable Disease Control and Prevention Communicable Disease Control Unit San Francisco, CA 94102 Phone: (415) 554-2830 Fax: (415) 554-2848 www.sfdph.org/cdcp

Measles: Frequently Asked Questions

Is measles serious?

For most people who get measles, the illness is not serious. It starts with fever, cough, runny nose, and watery eyes. After a few days a rash develops and lasts about a week. Then, it gets better by itself. However for some people who get measles, the disease can be more serious.

What happens when the disease is more serious?

Measles can infect other parts of the body. About 8 per 100 people with measles get diarrhea (8%), 7 per 100 get a middle ear infection (mainly young children), and 6 per 100 get pneumonia (lung infection with breathing trouble).

About 1 in 1,000 people with measles get encephalitis, a serious brain infection. Measles illness during pregnancy can cause early labor, miscarriage, and low birth weight infants. Measles in people with AIDS or weak immune systems can be very severe. In the U.S. people can still die from measles (about 2 per 1,000, usually related to pneumonia or encephalitis).

Is there treatment for measles?

There is no medicine that kills the measles virus once someone develops measles. Most people with measles get better by themselves. They should rest, drink plenty of fluids, and can take acetaminophen (Tylenol) to help reduce the fever and feel more comfortable.

People who are severely ill or who develop pneumonia, middle ear infection, or diarrhea should discuss treatment with their doctor. Although no medicine will make the measles virus go away, there may be a role for other medicines, such as antibiotics, in treating other possible reasons for infection.

How is measles spread or passed to others?

Measles is spread by bodily fluids – drops of saliva from the mouth, mucus from the nose, coughing or sneezing, and tears from the eyes.

The measles virus can also float in the air for up to 2 hours after someone with measles coughs or sneezes indoors. Someone who breathes air containing the measles virus can get measles.

People with measles can spread the disease to others from 4 days before the rash starts (before they know they have the disease), through 4 days after the rash begins.

What does it mean to be "exposed" to measles?

"Exposed" means you've been in a situation where you can catch the virus from someone who already has measles. For example, you can be exposed to the measles virus by being in the same room, home,

San Francisco Department of Public Health



Gavin Newsom Mayor Communicable Disease Control and Prevention Communicable Disease Control Unit San Francisco, CA 94102 Phone: (415) 554-2830 Fax: (415) 554-2848 www.sfdph.org/cdcp

office, or waiting room with someone who has measles or by being there up to 2 hours after the person with measles has left.

Is it true that some people can't get measles?

Yes. Some people are <u>immune</u> to measles, meaning that their body has already learned how to fight off the virus, and they won't become sick from it.

People can become immune to measles in 2 ways. If they've already had measles earlier in life, then they are immune and won't get it again. Or, if they have gotten 2 doses of measles vaccine, they should be immune (protected from getting measles).

How do you know if someone is immune to measles?

The only way to tell for sure that a person is protected (immune) is by a blood test. The blood test shows whether the body has antibodies to fight off the virus. This is called laboratory evidence of immunity.

Even if someone hasn't had the blood test, they are still considered immune if:

- 1. they had measles in the past, and have medical records to prove it, OR
- 2. they <u>received 2 doses of measles-containing vaccine</u> in the past (often given as MMR [measlesmumps-rubella] vaccine) and have medical records to prove it, OR
- 3. They were born before 1957

ABOUT MEASLES VACCINE

How well does the measles vaccine work?

The measles vaccine works extremely well. If 1,000 people each get 2 doses of measles vaccine, about 997 of them will be protected against measles (immune).

Getting 2 doses of vaccine is important, because some people (about 2-5 per 100) do not become protected (immune) after just one dose of vaccine.

Does the measles vaccine protect against other diseases too?

In the USA, measles vaccine is almost always given as a combination with vaccine against 2 other viruses: mumps and rubella. This combination vaccine is called MMR (measles, mumps, and rubella) and it gives protection against all 3 viruses.

What are the risks from MMR (measles, mumps, and rubella) vaccine?

A vaccine, like any medicine, is capable of causing serious problems, such as severe allergic reactions. The risk of MMR vaccine causing serious harm, or death, is extremely small. Getting MMR vaccine is much safer than getting any of these three diseases. Most people who get MMR vaccine do not have any problems with it.

City and County of San Francisco

San Francisco Department of Public Health



Gavin Newsom Mayor Communicable Disease Control and Prevention Communicable Disease Control Unit San Francisco, CA 94102 Phone: (415) 554-2830 Fax: (415) 554-2848 www.sfdph.org/cdcp

Mild Problems

- Fever (up to 1 person out of 6)
- Mild rash (about 1 person out of 20)
- Swelling of glands in the cheeks or neck (rare)

Moderate Problems

- Seizure (jerking or staring) caused by fever (about 1 out of 3,000 doses)
- Temporary pain and stiffness in the joints, mostly in teenage or adult women (up to 1 out of 4)
- Temporary low platelet count, which can cause a bleeding disorder (about 1 out of 30,000 doses)

Severe Problems (Very Rare)

• Serious allergic reaction (less than 1 out of a million doses)

What is Immunoglobulin (IG) and who needs it?

IG is a medical way to prevent measles from becoming more serious in some people. It will be given to some infants <12 months of age, pregnant women, persons with weakened immune systems, and others who for medical reasons can not get the vaccine.

What are the risks of IG?

Some people who get IG experience local discomfort, and some experience pain at the site of administration. Less common reactions include flushing, headache, chills and nausea. Serious reactions are uncommon. They may involve chest pain, difficulty breathing, anaphylaxis, low blood pressure, or shock. Health care professionals who administer IG should have training and resources to treat these rare but serious reactions.

ABOUT QUARANTINE

Who needs to be quarantined?

People who have been exposed to someone with measles and are in the period where they could still get sick from it.

Why is the quarantine period for measles 18 days?

Although the waiting period is usually 10-14 days, sometimes it can take longer from the time someone is exposed to measles until the time they get sick from it.

If I get the vaccine now, will the quarantine be shorter?

No. But getting vaccine now may prevent you from getting sick.

San Francisco Department of Public Health



Gavin Newsom Mayor Communicable Disease Control and Prevention Communicable Disease Control Unit San Francisco, CA 94102 Phone: (415) 554-2830 Fax: (415) 554-2848 www.sfdph.org/cdcp

Do household members of a quarantined person need to be vaccinated?

See above: "How do you know if someone is immune to measles?" Household members who are immune to measles do not need vaccine. Household members who are not immune, or are not sure whether they are immune, should consider getting vaccine.

Can anyone come into the house where a person is quarantined?

No. Entering the home where a person is quarantined is highly discouraged. However, in cases requiring caregivers such as a child or the elderly, some individuals will be allowed to enter the household.

Documentation of birth before 1957, two valid measles vaccine doses, or a blood test showing measles immunity will need to be seen by Public Health officials for persons wanting to enter the quarantined home. No others are allowed to enter.

What about handwashing? Wearing a mask?

It is important for people to wash their hands frequently with soap and water, especially after coughing, sneezing, or blowing their nose, or after going to the bathroom, or having contact with moist materials such as tissues, pampers, and used masks. A minimum of 60% alcohol-based hand sanitizer can also be used instead of soap and water if the hands are not visibly dirty.

If there are other household members living in the home of the quarantined person, and who are not immune to measles, the likelihood that they will get sick with measles may be reduced if the quarantined person stays inside their designated room/area, avoids contact with others, and wears a mask when in common rooms. If the quarantined person cannot wear a mask, household members could wear one when they have contact with the quarantined individual.

Does the quarantined person need to wear a mask when leaving the house?

The quarantined person should not leave the house, except to receive emergency medical care. In this case, yes, the quarantined person should wear a mask. If the quarantined person will not be able to wear the mask at all times because they are a baby or a child, this should be discussed with the medical facility before arrival.

Can pets get infected with measles or spread measles?

No, pets do not get infected with or spread the measles virus.