

Test Order

Shiga Toxin Producing Escherichia coli (STEC) Shiga Toxin Gene PCR

Synonym(s)	STEC PCR, stx1/stx2 gene PCR, Shiga Toxin Gene PCR
Methodology	Real Time Shiga Toxin Gene PCR
Acceptable Specimen	Pure isolates, previously identified as <i>Escherichia coli</i> ; GN or Mac broth;
Type(s) for Testing	Stool in PARA-PAK C&S.
Transport / Collection	Pure isolates: transport on nutrient agar slant, packaged and sealed
Medium	according to regulations. PARA-PAK C&S: mix well after collection.
	Refrigerate, and deliver to the laboratory within 3 days of collection. GN/Mac
	broths: packaged and sealed according to regulations.
Storage and	Refrigerate until transported to the laboratory. Transport at room
Preservation of	temperature.
Specimen	
Minimum Volume	PARA-PAK C&S: Fill to line (do NOT overfill).
Required	
Additional Collection	N/A
Instructions	
Additional Required	N/A
Information	
Send Out?	No
Turnaround Time	10 business days (M-F)
Testing Restrictions	
Requisition Form(s)	http://sfcdcp.org/document.html?id=1036
Limitations / Notes /	Delays in GN/Mac broth submissions to the laboratory can lead to reduced
Disclaimers	success in STEC isolation.
	This PCR assay was developed and its performance characteristics were
	determined by the San Francisco Public Health Laboratory. It has not been
	cleared or approved by the U.S. Food and Drug Administration. The FDA has
	determined that such approval is not necessary and results should not be
	regarded as investigational or for research.

Updated 10/03/2022